



EUSPC06JUN2012



HUMULIN® R
REGULAR
INSULIN HUMAN INJECTION, USP (rDNA ORIGIN)
100 UNITS PER ML (U-100)

HUMULIN® N
NPH
HUMAN INSULIN (rDNA ORIGIN) ISOPHANE SUSPENSION
100 UNITS PER ML (U-100)

HUMULIN® 70/30
70% HUMAN INSULIN ISOPHANE SUSPENSION
AND 30% HUMAN INSULIN INJECTION (rDNA ORIGIN)
100 UNITS PER ML (U-100)

1. NAME OF THE MEDICINAL PRODUCT

Humulin R 100 IU/ml solution for injection in vial
Humulin N 100 IU/ml suspension for injection in vial
Humulin 70/30 100 IU/ml suspension for injection in vial

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 ml contains 100 IU human insulin (produced in *E.coli* by recombinant DNA technology).
Humulin R
One vial contains 10 ml equivalent to 1000 IU of soluble insulin.

Humulin N
One vial contains 10 ml equivalent to 1000 IU of isophane insulin.

Humulin 70/30
One vial contains 10 ml equivalent to 1000 IU of biphasic isophane insulin –
30 % soluble insulin / 70 % isophane insulin.

For a full list of excipients, see section 6.1.

Not all strengths or presentations may be marketed.

3. PHARMACEUTICAL FORM

Humulin R

A solution for injection in a vial.
Humulin R is a sterile, clear, colourless, aqueous solution of human insulin.

Humulin N

A suspension for injection in a vial.
Humulin N is a sterile suspension of a white, crystalline precipitate of isophane human insulin in an isotonic phosphate buffer.

Humulin 70/30

A suspension for injection in a vial.
Humulin 70/30 is a sterile suspension of human insulin in the proportion of 30 % soluble insulin to 70 % isophane insulin.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

For the treatment of patients with diabetes mellitus who require insulin for the maintenance of glucose homeostasis.

4.2 Posology and method of administration

The dosage should be determined by the physician, according to the requirement of the patient.

Subcutaneous administration should be in the upper arms, thighs, buttocks or abdomen. Use of injection sites should be rotated so that the same site is not used more than approximately once a month.

Care should be taken when injecting any Humulin insulin preparations to ensure that a blood vessel has not been entered. After any insulin injection, the injection site should not be massaged. Patients must be educated to use proper injection techniques.

Each pack contains a patient information leaflet with instructions on how to inject insulin.

Humulin R

Humulin R should be given by subcutaneous injection but may, although not recommended, also be given by intramuscular injection. It may also be administered intravenously.

Humulin N

Humulin N should be given by subcutaneous injection but may, although not recommended, also be given by intramuscular injection. This formulation should not be administered intravenously.

Humulin N may be administered in combination with Humulin R (See Instructions for use and handling - for Mixing of Insulins).

Humulin 70/30

Humulin 70/30 should be given by subcutaneous injection but may, although not recommended, also be given by intramuscular injection. This formulation should not be administered intravenously.

Humulin Mixture formulation is a ready-made defined mixture of soluble and isophane insulin designed to avoid the need for the patient to mix insulin preparations. A patient's treatment regimen should be based on their individual metabolic requirements.

4.3 Contraindications

Hypoglycaemia.

Hypersensitivity to Humulin or to the formulation excipients, unless used as part of a desensitisation programme.

Under no circumstances should any Humulin formulation other than Humulin R be given intravenously.

4.4 Special warnings and precautions for use

Transferring a patient to another type or brand of insulin should be done under strict medical supervision. Changes in strength, brand (manufacturer), type (soluble, isophane, mixture), species (animal, human, human insulin analogue), and/or method of manufacture (recombinant DNA versus animal-source insulin) may result in the need for a change in dosage.

Some patients taking human insulin may require a change in dosage from that used with animal-source insulins. If an adjustment is needed, it may occur with the first dose or during the first several weeks or months.

A few patients who experienced hypoglycaemic reactions after transfer to human insulin have reported that the early warning symptoms were less pronounced or different from those experienced with their previous animal insulin. Patients whose blood glucose is greatly improved, e.g. by intensified insulin therapy, may lose some or all of the warning symptoms of hypoglycaemia and should be advised accordingly. Other conditions which may make the early warning symptoms of hypoglycaemia different or less pronounced include long duration of diabetes, diabetic nerve disease, or medications such as beta blockers. Uncorrected hypoglycaemic and hyperglycaemic reactions can cause loss of consciousness, coma or death.

The use of dosages which are inadequate or discontinuation of treatment, especially in insulin-dependent diabetics, may lead to hyperglycaemia and diabetic ketoacidosis; conditions which are potentially lethal.

Treatment with human insulin may cause formation of antibodies, but titres of antibodies are lower than those to purified animal insulin.

Insulin requirements may change significantly in diseases of the adrenal, pituitary or thyroid glands and in the presence of renal or hepatic impairment.

Insulin requirements may be increased during illness or emotional disturbances.

Adjustment of insulin dosage may also be necessary if patients change their level of physical activity or change their usual diet.

Combination of human insulin with pioglitazone

Cases of cardiac failure have been reported when pioglitazone was used in combination with insulin, especially in patients with risk factors for development of cardiac heart failure. This should be kept in mind, if treatment with the combination of pioglitazone and human insulin is considered. If the combination is used, patients should be observed for signs and symptoms of heart failure, weight gain and oedema. Pioglitazone should be discontinued, if any deterioration in cardiac symptoms occurs.

4.5 Interactions with other medicinal products and other forms of interaction

A number of medicinal products are known to interact with glucose metabolism and therefore the physician should be consulted when using other medications in addition to human insulin (see section 4.4). The physician must therefore take possible interactions into account and should always ask his patients about any medicinal products they take.

Insulin requirements may be increased by substances with hyperglycaemic activity, such as glucocorticoids, thyroid hormones, growth hormone, danazol, beta₂-sympatomimetics (such as ritodrine, salbutamol, terbutaline), thiazides.

Insulin requirements may be reduced in the presence of substances with hypoglycaemic activity, such as oral hypoglycaemics (OHA), salicylates (for example, acetylsalicylic acid), certain antidepressants (monoamine oxidase inhibitors), certain angiotensin converting enzyme (ACE) inhibitors (captopril, enalapril), angiotensin II receptor blockers, non-selective beta-blocking agents and alcohol.

Somatostatin analogues (octreotide, lanreotide) may both decrease or increase insulin dose requirements.

4.6 Fertility, pregnancy and lactation

It is essential to maintain good control of the insulin treated (insulin-dependent or gestational diabetes) patient throughout pregnancy. Insulin requirements usually fall during the first trimester and increase during the second and third trimesters. Patients with diabetes should be advised to inform their doctors if they are pregnant or are contemplating pregnancy.

Careful monitoring of glucose control, as well as general health, is essential in pregnant patients with diabetes.

Patients with diabetes who are lactating may require adjustments in insulin dose and/or diet.

4.7 Effects on ability to drive and use machines

The patient's ability to concentrate and react may be impaired as a result of hypoglycaemia. This may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery).

Patients should be advised to take precautions to avoid hypoglycaemia whilst driving, this is particularly important in those who have reduced or absent awareness of the warning signs of hypoglycaemia or have frequent episodes of hypoglycaemia. The advisability of driving should be considered in these circumstances.

4.8 Undesirable effects

Hypoglycaemia is the most frequent undesirable effect of insulin therapy that a patient with diabetes may suffer. Severe hypoglycaemia may lead to loss of consciousness, and in extreme cases, death. No specific frequency for hypoglycaemia is presented, since hypoglycaemia is a result of both the insulin dose and other factors e.g. a patient's level of diet and exercise.

Local allergy in patients is common (1/100 to < 1/10). Redness, swelling, and itching can occur at the site of insulin injection. This condition usually resolves in a few days to a few weeks. In some instances, local reactions may be related to factors other than insulin, such as irritants in the skin cleansing agent or poor injection technique.

Systemic allergy, which is very rare (< 1/10,000) but potentially more serious, is a generalised allergy to insulin. It may cause rash over the whole body, shortness of breath, wheezing, reduction in blood pressure, fast pulse, or sweating. Severe cases of generalised allergy may be life-threatening.

In the rare event of a severe allergy to Humulin, treatment is required immediately. A change of insulin or desensitisation may be required.

Lipodystrophy at the injection site is uncommon (1/1,000 to < 1/100).

Cases of oedema have been reported with insulin therapy, particularly if previous poor metabolic control is improved by intensified insulin therapy.

4.9 Overdose

Insulin has no specific overdose definitions, because serum glucose concentrations are a result of complex interactions between insulin levels, glucose availability and other metabolic processes. Hypoglycaemia may occur as a result of an excess of insulin relative to food intake and energy expenditure.

Hypoglycaemia may be associated with listlessness, confusion, palpitations, headache, sweating and vomiting.

Mild hypoglycaemic episodes will respond to oral administration of glucose or sugar products.

Correction of moderately severe hypoglycaemia can be accomplished by intramuscular or subcutaneous administration of glucagon, followed by oral carbohydrate when the patient recovers sufficiently. Patients who fail to respond to glucagon must be given glucose solution intravenously.



carbohydrate when the patient recovers sufficiently. Patients who fail to respond to glucagon must be given glucose solution intravenously.



PA909SPXS00

PA909SPXS00

PA909SPXS00

PA909SPXS00

Item Code PA909SPXS00		Previous Item Code (to be destroyed) N/A			
PPD Information Box				ALRP Information Box	
Start Date 02 DEC 2014	Proof No.: 1	Printing Colours 1/1 BLACK	Technical Colours Die Cut	Affiliate Barcode: Type: N/A Code: N/A	Translations of Variable Data
Technical Information: Layout name ALCPA0050A00	Size (mm): 110 x 592			Other Regulated Elements N/A	lot: N/A
	Folded Size (mm) 110x296				mfq date: N/A
BLUE Project No. 1400000375	No. of Pages: 1/2				exp date: N/A
	Sick Code N/A				price: N/A
Feed Direction: (For labels only)		<input checked="" type="checkbox"/> N/A <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>			
				Site: Alcobendas	
		Riverside Drive, Cleckheaton, West Yorkshire, BD19 4DH, United Kingdom Tel: +44 (0) 127 484 8000 www.mbs.co.uk		Internal to MBS (no lilly check required) MBS Artwork Ref: 350533 Operator Name: JL	

If the patient is comatose, glucagon should be administered intramuscularly or subcutaneously. However, glucose solution must be given intravenously, if glucagon is not available or if the patient fails to respond to glucagon. The patient should be given a meal as soon as consciousness is recovered.

Sustained carbohydrate intake and observation may be necessary because hypoglycaemia may occur after apparent clinical recovery.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Humulin R

Pharmaco-therapeutic group: Humulin R: ATC code A10A B01.
Humulin R is a rapidly acting insulin preparation.

Humulin N

Pharmaco-therapeutic group: Humulin N: ATC code A10A C01.
Humulin N is an intermediate acting insulin preparation.

Humulin 70/30


Pharmaco-therapeutic group: Humulin 70/30: ATC code A10A D01.
Humulin 70/30 is an intermediate acting insulin preparation.

The prime activity of insulin is the regulation of glucose metabolism.


In addition insulin has several anabolic and anti-catabolic actions on a variety of different tissues. Within muscle tissue this includes increasing glycogen, fatty acid, glycerol and protein synthesis and amino acid uptake, while decreasing glycogenolysis, gluconeogenesis, ketogenesis, lipolysis, protein catabolism and amino acid output.

The typical activity profile (glucose utilisation curve) following subcutaneous injection is illustrated below by the heavy line. Variations that a patient may experience in timing and/or intensity of insulin activity are illustrated by the shaded area. Individual variability will depend on factors such as size of dose, site of injection temperature and physical activity of the patient.


Humulin R



Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30



Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N



Humulin 70/30



Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N



Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N




Humulin 70/30




Humulin R




Humulin N



Humulin 70/30



Humulin R



Humulin N



Humulin 70/30



Humulin R



Humulin N



Humulin 70/30




Humulin R



Humulin N



Humulin 70/30




Humulin R



Humulin N



Humulin 70/30



Humulin R



Humulin N




Humulin 70/30




Humulin R



Humulin N



Humulin 70/30



Humulin R



Humulin N

Humulin 70/30

Humulin R

Humulin N

Humulin 70/30

Humulin R

Humulin N

Humulin 70/30

